Volume 2, Issue 1, January 2025

Network Structure and Efficiency Impact on Graph Labeling

Amar L. Powar Shivaji University, Kolhapur, India

Abstract: In this paper is proposed to Cordial labeling and vertex n-magic labeling that two types of methods. Using this cordial labeling is best one because vertex n-magic labeling is used 1 to n value but the cordial labeling is used 0 and 1 only. So, the paper assign 0 and 1 is the best one.

Keywords: Cordial labeling, binary vertex labeling, apex vertex, pendant vertex

I. INTRODUCTION

A graph is cordial if it is possible to label its vertices with 0's and 1's so that when the edges are labeled with the difference of the labels at their endpoints, the number of vertices labeled with ones and zeros differ at most by one. An interesting vertex labeling with numbers is vertex n-magic. Vertex n-magic graphs are graphs labeled with numbers in which every vertex and its incident edges add up to the same number. This number is called the magic number. It is still does not shows that what types of graphs are vertex-magic and which are not. The cycle graph is one type of graph that has interesting vertex-magic properties. The degree of a vertex, denoted f(v) in a graph is the number of edges incident to it. A vertex with degree zero is called an isolated vertex. That is, a vertex which is not an endpoint of any edge. A vertex with degree one is called leaf vertex. A vertex is said to be an apex vertex, if all the chords of the vertices are concurrent.

Definition 1. Cordial labeling

Let G = (V,E) be a simple graph, and f be the function from $V(G) \rightarrow (0,1)$ and for edge uv assign the label

f(u) - f(v). f is called a cordial labeling.

Copyright to JSRD www.banglajo.in

Volume 2, Issue 1, January 2025



Definition 3. Binary vertex labeling A mapping $f : V(G) \{0,1\}$ is called binary vertex labelling of G and f(V) is called the label of the vertex V of G under f.

Definition 4. Even vertex magic total graph A labeling is said to be an even vertex N-magic total if $f(V(G)) = \{2,4,6,\ldots,\}$. A graph that admits an even vertex N-magic total labeling is called even vertex magic total graph.

Definition 5. Shadow graph The connected graph G is said to be a Shadow graph, it is constructed by taking two copies of G say G' and G". Join each vertex u' in G' to the neighbours of the corresponding vertex v' in G".

Definition 6. Splitting graph A graph G is said to be the Splitting graph S'(G) of a graph G is obtained a new vertex V' corresponding to each vertex V of G such that N(V)=N(V'). S'(k1,n) is cordial graph.

Volume 2, Issue 1, January 2025

Proof:

Let v_1, v_2, \ldots, v_n be the pendant vertices and v be the apex vertex of $k_{1,n}$ and u_1, u_2, \ldots, u_n are added vertices corresponding to v_1, v_2, \ldots, v_n to obtain S^{*}(k_1, n).

Let G be the Splitting graph S'(k1,n) then

```
|V(G)| = 2n + 2 and
```

|E(G)| = 3n.

The labeling is a function from $f: V(G) \rightarrow \{0,1\}$ is defined as per the following cases

Case (i)

n

$\equiv 0 \pmod{6}$	n = 6,12,18,
f(u) = 1,	
$f(u_i)=0;if\;i\equiv 1,3,(mod\;6)$	i = 7,9,
= 1; if $i \equiv 0, 2 \pmod{6}$	i = 6,8,
f(v) = 0,	
$f(v_i) = 0; \text{ if } i \equiv 3, 5 \pmod{6}$	i = 9.11,
$= 1; $ if $i \equiv 2, 4 \pmod{6}$	i = 8,10,

Case (ii)

```
n \equiv 1 \pmod{8} n=9,17,25,...
```

f(u) = 1

 $f(u_i) = 0$; if $i \equiv 5, 7 \pmod{8}$ i = 13, 15, ...,

 $=1: \text{ if } i \equiv 4, 6 \pmod{8}$ $i = 12, 14, \dots$

$$f(v) = 0,$$

$$f(v_i) = 0$$
; if $i \equiv 1, 3 \pmod{8}$ $i = 17,19$

$$=1$$
; if $i \equiv 0, 2 \pmod{8}$ $i = 16, 18, ...$

Case (iii)

Copyright to JSRD www.banglajo.in

Volume 2, Issue 1, January 2025

$n \equiv 2,3 \pmod{8}$	n=10,11,
f(u) = 1,	
$f(u_i) = 0; if i \equiv 0, 1 \; (mod \; 8)$	$i = 8, 9, \ldots$
$-1; if i \equiv 2, 3 \pmod{8}$	i = 10, 11,
f(v) = 0;	
$f(v_i)-0; if \; i\equiv 2,3 \; (mod \; 8)$	$i = 18, 19, \ldots$
-1 ; if $i \equiv 0, 1 \pmod{8}$	i – 16,17,

The condition satisfies the labeling pattern in all the above cases.

 $\left\| v_f(0) - v_f(1) \right\| \le 1$ and $\left\| e_f(0) - e_f(1) \right\| \le 1$

in each case which is shown in table.

Hence, S'(k1,n) is cordial graph

Let n - 4a + b, where $n \in N$

Table

b	Vertex condition	Edge condition
0,2	$v_{f}(0) = v_{f}(1)$	$e_{f}(0) = e_{f}(1)$
1	$v_{f}(0) = v_{f}(1)$	$e_f(0) = e_f(1) + 1$
3	$v_{f}(0) = v_{f}(1)$	$e_f(0) + 1 = e_f(1)$

Example



Copyright to JSRD www.banglajo.in

Volume 2, Issue 1, January 2025

Theorem.2

For each m, n \ge 3 and m, n odd, there exists an even vertex magic total labeling of $C_m \times C_n$ with the magic constant $k = \frac{17}{2} \text{ mn} + \frac{5}{2}$.

Proof.

Let $G \cong C_m \times C_n$ have vertices $v_{i,j}$ vertical edges $v_{i,j} v_{i+1,j}$ and horizontal edges $v_{i,j} v_{i,j+1}$

where $i = 0, 1, \dots, m - 1$,

 $j = 0, 1, \dots, n - 1$ and m and n are odd integers greater than 1.

Let us consider the following labeling, where i and j are the subscripts taken modulo m and n respectively.

$$f(v_{i,i}) = 2 (mn - jm - i)$$

$$f\left(v_{i,j}v_{i+1,j}\right) = \begin{cases} 2m(n+j+1) - 1 - i & \text{ if i is even} \\ \\ 2m(n+j) + m - i - 1 & \text{ if i is odd} \end{cases}$$

$$\begin{split} f(v_{i,j}v_{i,j+1}) = & \begin{cases} 2(mn-m+i)-jm+1 & \text{ if } j \text{ is even} \\ 2(i-m)-jm+mn+1 & \text{ if } j \text{ is odd} \end{cases}$$

Volume 2, Issue 1, January 2025

Case (i)

If both i and j are even then the magic constant k is given by

$$\begin{aligned} & \text{When } i=2, \, j=2 \\ & \text{ } k \ = \ f\left(v_{i,j}\right) \ + \ f\left(v_{i-1,j}v_{i,j}\right) \ + \ f\left(v_{i,j},v_{i+1,j}\right) \ + \ f\left(v_{i,j-1}v_{i,j}\right) \ + \ f\left(v_{i,j},v_{i,j+1}\right) \end{aligned}$$

$$k = f(v_{2,2}) + f(v_{2-1,2}v_{2,2}) + f(v_{2,2},v_{2+1,2}) + f(v_{2,2-1}v_{2,2}) + f(v_{2,2},v_{2,2+1})$$

$$= 2(mn - 2m - 2) + 2m(n + 2) + m - (2 - 1) - 1 + 2m(n + 2 + 1)$$

-1 - 2 + 2(2 - m) - (2 - 1)m + mn + 1 + 2(mn - m + 2) - 2m + 1

k = 9mn + 1

Case (ii)

If i is even and j is odd then the magic constant k is given by

The magic constants are same in all the cases and. $f(V(G)) = \{2, 4, ..., 2p\}$.

Therefore G is even vertex magic total with the magic constant k = 9mn + 1.

Copyright to JSRD www.banglajo.in

Volume 2, Issue 1, January 2025



II. CONCLUSION

From this, we conclude labeling cordial is most excellent private since pinnacle n-magic labeling is worn 1 to n worth but the affable labeling is worn 0 and 1 simply. So, the rag consign 0 and 1 is the finest solitary.

REFERENCES

[1] Amit H .Rokad, Kalpesh M. Patadiya "Cordial labeling of some graphs". Aryabhatta Journal of Mathematics and Informatics. (2017).

[2] Marimuthu and G.Kumar "Vertex N-magic total labeling of graphs". (2016).

[3] J.A.Bondy, and U.S.R. Murty, Graph theory with Applications.North Holland, Newyork.